

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 9/127, 31/71</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/12155 (43) International Publication Date: 9 June 1994 (09.06.94)</p>
<p>(21) International Application Number: PCT/US92/10591 (22) International Filing Date: 2 December 1992 (02.12.92)</p> <p>(71) Applicant (for all designated States except US): VESTAR, INC. [US/US]; 650 Cliffside Drive, San Dimas, CA 91773 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): PROFFITT, Richard, T. [US/US]; 11 N. Altura Road, Arcadia, CA 91007 (US). BRACKEN, Kevin, R. [US/US]; 8226 Day Street, Sunland, CA 91040 (US). CHIANG, Su-Ming [US/US]; 22009 Napa Street, Canoga Park, CA 91304 (US).</p> <p>(74) Agent: COCHRAN, Adam; Vestar, Inc., 650 Cliffside Drive, San Dimas, CA 91773 (US).</p>		<p>(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>

(54) Title: ANTIBIOTIC FORMULATION AND PROCESS

(57) Abstract

A liposomal aminoglycoside formulation comprising a neutral lipid, a negatively charged lipid and a sterol. The formulation contains unilamellar vesicles having an average size below 100 nm. A process of making liposomes containing an aminoglycoside is provided where the hydration temperature is significantly below the transition temperature of the formulation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ANTIBIOTIC FORMULATION AND PROCESS

Field of the Invention

This invention relates to the fields of biochemistry and medicine, and particularly to a liposome formulation. More specifically, it relates to a liposomal formulation containing an aminoglycoside, its process of manufacture and its use. This invention also relates to formulations having reduced toxicity, longer stability, and superior efficacy.

Background of the Invention

The discovery of aminoglycosides began in the 1940s with the isolation of streptomycin from *Streptomyces griseus*. Since the 1940s, other aminoglycosides have been discovered and synthesized. These include neomycin which is obtained or isolated from *Streptomyces fradiae*; kanamycin which is isolated from *Streptomyces kanamyceticus*; gentamicin which is isolated from *Micromonospora purpurea*; tobramycin which is isolated from *Streptomyces tenebrarius*; sisomicin isolated from *micromonospora inyoensis*; amikacin which is a semisynthetic derivative of Kanamycin A; and netilmicin which is a semisynthetic derivative of sisomicin. Amikacin has the broadest spectrum of antimicrobial activity of all the aminoglycosides. It also has a unique resistance to the immunoglycoside-inactivating enzymes.

The aminoglycosides are polar-cations which consist of two or more amino sugars joined in a glycosidic linkage to a hexose nucleus, which is usually in a central position. The aminoglycosides are used primarily to treat infections caused by gram-negative bacteria. However, aminoglycosides have been used in recent years to treat bacteria from the genera *Mycobacteria*. For example, amikacin has shown to be effective against *Mycobacterium tuberculosis*. Aminoglycosides have also been tested against *M. avium* infections including *M. avium-intracellulare* complex (MAC) which is a group of related acid-fast organisms that grow only slightly faster than *M. tuberculosis* and can be divided into a number of serotypes. At the beginning of the twentieth century, tuberculosis was the most prevalent cause of death in the United States. By the

late 1940s, with the advent of streptomycin, tuberculosis infection had decreased substantially. Since the mid-1980s with the appearance of the acquired immune deficiency syndrome, tuberculosis again began to emerge as a major health problem. Further, the new cases of tuberculosis showed resistance to many of the available antibiotic therapies. Similarly MAC, once considered rare, is now the most common systemic bacterial type infections in patients suffering from acquired immune deficiency syndrome. Hence, the search for an effective antibiotic has intensified.

Although the aminoglycosides have been useful in treating infections, the use of these antibiotics is not free from toxicity and side effects. Aminoglycosides may produce irreversible vestibular, cochlear, and renal toxicity. The two main toxic effects of aminoglycosides are ototoxicity and nephrotoxicity. Studies have found that the aminoglycosides antibiotics may cause polyuria, decreased urinary osmolality, proteinuria, enzymuria, glycosuria, and a decrease in the rate of glomerular filtration. Some investigators believe that nephrotoxicity results from the accumulation of the aminoglycosides in the renal cortex because of the long half-life of the agents in that tissue.

Liposomes are microscopic vesicles made, in part, from phospholipids which form closed, fluid filled spheres when dispersed with water. Phospholipid molecules are polar, having a hydrophilic ionizable head and two hydrophobic tails consisting of long fatty acid chains. Thus, when sufficient phospholipid molecules are present with water, the tails spontaneously associate to exclude water while the hydrophilic phosphate heads interact with water. The result is a bilayer membrane in which the fatty acid tails converge in the newly formed membrane's interior and the polar heads point in opposite directions toward an aqueous medium. These bilayer membranes can be caused to form closed spheres known as liposomes. The polar heads at the inner surface of the membrane point toward the aqueous interior of the liposome. At the opposite surface of the spherical membrane, the polar heads interact with the surrounding aqueous medium. As the liposomes are formed, water soluble molecules can be

incorporated into the aqueous interior, and lipophilic molecules will tend to be incorporated into the lipid bilayer. Liposomes may be either multilamellar, like an onion with liquid separating many lipid bilayers, or unilamellar, with a single bilayer surrounding an entirely liquid center.

5 There are many types of liposome preparation techniques which may be employed and which produce various types of liposomes. These can be selected depending on the use, the chemical intended to be entrapped, and the type of lipids used to form the bilayer membrane. The requirements which must be considered in producing a liposome preparation are similar to those of other
10 controlled release mechanisms. They are as follows: (1) high percent of chemical entrapment; (2) increased chemical stability; (3) low chemical toxicity; (4) rapid method of production; and (5) reproducible size distribution.

 The first method described to encapsulate chemicals in liposomes involved production of multilamellar vesicles (MLVs). Methods for encapsulating
15 chemicals in MLVs are known in the art.

Liposomes can also be formed as unilamellar vesicles (UVs), which generally have a size less than 1 μm . There are several techniques known in the art which are used to produce unilamellar liposomes.

 Smaller unilamellar vesicles can be formed using a variety of techniques.
20 By dissolving phospholipids in ethanol and injecting them into a buffer, the lipids will spontaneously rearrange into unilamellar vesicles. This provides a simple method to produce UVs which have internal volumes similar to that of those produced by sonication (0.2-0.5 L/mol/lipid). Sonication or extrusion (through filters) of MLVs also results in dispersions of UVs having diameters of up to 0.2
25 μm , which appear as clear or translucent suspensions.

 Another common method for producing small UVs is the detergent removal technique. Phospholipids are solubilized in either ionic or non-ionic detergents such as cholates, Triton X, or n-alkylglucosides. The drug is then mixed with the solubilized lipid-detergent micelles. Detergent is then removed
30 by one of several techniques: dialysis, gel filtration, affinity chromatography,

centrifugation, ultrafiltration. The size distribution and entrapment efficiencies of the UVs produced this way will vary depending on the details of the technique used.

The therapeutic use of liposomes includes the delivery of drugs which are normally toxic in the free form. In the liposomal form the toxic drug may be directed away from the sensitive tissue and targeted to selected areas. Liposomes can also be used therapeutically to release drugs, over a prolonged period of time, reducing the frequency of administration. In addition, liposomes can provide a method for forming an aqueous dispersion of hydrophobic drugs for intravenous delivery.

When liposomes are used to target encapsulated drugs to selected host tissues, and away from sensitive tissues, several techniques can be employed. These procedures involve manipulating the size of the liposomes, their net surface charge as well as the route of administration. More specific manipulations have included labeling the liposomes with receptors or antibodies for particular sites in the body.

The route of delivery of liposomes can also affect their distribution in the body. Passive delivery of liposomes involves the use of various routes of administration, e.g., intravenous, subcutaneous and topical. Each route produces differences in localization of the liposomes. Two common methods used to actively direct the liposomes to selected target areas are binding either antibodies or specific receptor ligands to the surface of the liposomes. Antibodies are known to have a high specificity for their corresponding antigen and have been shown to be capable of being bound to the surface of liposomes, thus increasing the target specificity of the liposome encapsulated drug.

Since the chemical composition of many drugs precludes their intravenous administration, liposomes can be very useful in adapting these drugs for intravenous delivery. Many hydrophobic drugs, including cyclosporine, fall into this category because they cannot be easily dissolved in a water-based medium and must be dissolved in alcohols or surfactants which have been shown to cause

toxic reactions *in vivo*. Liposomes, composed of lipids, with or without cholesterol, are nontoxic. Furthermore, since liposomes are made up of amphipathic molecules, they can entrap hydrophilic drugs in their interior space and hydrophobic molecules in their lipid bilayer. Although methods for making liposomes are well known in the art, it is not always possible to determine a working formulation without undue experimentation.

Liposomal formulations containing aminoglycosides have been prepared. Many of the preparations include aerosol formulations using MLVs. Other formulations contain a large amount of negatively charged lipids, generally greater than 20%, to increase retention time or circulation half-life. Problems associated with aminoglycosides liposomal formulations include short retention time in the system because of RES uptake. Attempted formulations in the art have also resulted in liposomal aminoglycosides that are unstable - both on the shelf and in serum.

Thus, it is a desideratum to provide for a novel formulation which would increase the retention time of an aminoglycoside in a mammal's system and which can deliver more drug with superior efficacy and lower toxicity than free drug. It is also desirable to provide a process which allows the manufacture of a clear, stable and efficacious aminoglycoside liposome suspension because of the inability of those of ordinary skill to produce a therapeutically effective aminoglycoside unilamellar liposomal formulation having an average particle size of less than 100 nm, preferably with a high drug to total lipid ratio.

Summary of the Invention

Liposomes are provided in the present invention which comprise an aminoglycoside wherein the liposomes are unilamellar having an average size of 100 nm or less. A liposomal formulation is provided which comprises an aminoglycoside wherein the liposomes are comprised of a neutral lipid, a sterol and a negatively charged lipid. Small unilamellar vesicles are also provided wherein the molar amount of negatively charged lipid is less than 20% of total lipid. A preferred formulation is liposomes having an aminoglycoside wherein

the liposomes are unilamellar vesicles having an average size of 30 nm to 100 nm and further comprised of a phosphatidylcholine, cholesterol, and a phosphatidylglycerol wherein the molar amount of phosphatidylglycerol is about 3%. The drug to total lipid ratio is between 1:9 and 1:3 with the preferred ratio at about 1:4. A preferred formulation is also provided comprising liposomes including an aminoglycoside wherein the liposomes are unilamellar vesicles having an average size less than 100 nm wherein the lipids comprise hydrogenated soy phosphatidylcholine, cholesterol, and distearoylphosphatidylglycerol in a molar ratio of about 2:1:0.1.

10 The present invention also includes a method for making aminoglycoside liposomes. The process comprises forming a lipid powder comprised of a neutral phospholipid, a sterol, and a negatively charged lipid; mixing the powder with an aminoglycoside in an aqueous buffer and hydrating the mixture at a temperature significantly below the transition temperature of the lipid mixture; and reducing the size of the liposome to an average particle size of less than 100 nm.

The present invention also provides for the treatment of bacterial infections in mammals comprising preparing a liposomal formulation having an aminoglycoside wherein the liposomes are those described above and which are used to treat infections by introducing a therapeutic effective amount of the liposomes into a mammal. Thus, the present invention provides the use of liposomal aminoglycoside formulations to treat bacterial infections. The bacterial infections treated include opportunistic aerobic gram-negative bacilli such as the genera *Pseudomonas*. Another aspect of the invention includes the use of the liposomal formulations in the treatment of a bacterial infection caused by *P. aeruginosa*. The method of treating bacterial infections is not limited to gram-negative infections. The liposomal formulations can be used to treat bacterial infections comprising gram-positive bacilli such as the genera *Mycobacterium*. The invention is particularly useful for treating mycobacterium which causes tuberculosis-like diseases. Numerous bacterium may be treated using the

liposomes described above. The bacteria would include: *M. tuberculosis*, *M. leprae*, *M. Intracellulare*, *M. smegmatis*, *M. bovis*, *M. kansasii*, *M. avium*, *M. scrofulcium*, or *M. africanum*. Liposomal formulations of aminoglycoside are also particularly useful in treating MAC.

5 Many, if not all, mycobacterium infections discussed above are difficult to treat because the bacteria invade phagocytic cells such as macrophages. Application of liposomal formulations containing an aminoglycoside result in the intracellular delivery of the drug which would not normally occur with the delivery of the free drug. Thus, the present invention provides a treatment of
10 infected phagocytic cell in mammals by delivering a therapeutic or effective amount of an aminoglycoside into a macrophage using a unilamellar liposome having an average size of 100 nm or less.

Detailed Description of the Invention

As used herein, the term liposome refers to unilamellar vesicles or
15 multilamellar vesicles such as described in U.S. Patent 4,753,788 and 4,935,171, the contents of which are incorporated herein by reference.

The present invention provides liposomal aminoglycoside formulation preferably containing a phosphatidylcholine, a phosphatidylglycerol, cholesterol (CHOL) and amikacin. Hydrogenated soy phosphatidylcholine (HSPC),
20 distearoyl phosphatidylglycerol (DSPG) are the preferred lipids for use in the invention. Other suitable phosphatidylcholines include those obtained from egg or plant sources, or those that are partially or wholly synthetic. Other phosphatidylglycerols that may be used include dimyristoylphosphatidylglycerol (DMPG) and dilaurylphosphatidylglycerol (DLPG).

25 The preferred formulation includes liposomes comprising an encapsulated aminoglycoside wherein the liposomes are unilamellar vesicles having an average size of less than 100 nm and which are stable for at least two weeks at 22°C without a significant change in size or without loss of more than 10% of encapsulated aminoglycosides. The preferred ratio of HSPC:CHOL:DSPG is
30 about 2:1:0.1 and the drug to total lipid ratio is about 1:4. Other preferred

formulations include DSPG in a molar amount of 0 to 20%. Other preferred formulations include formulations where the drug to total lipid ratio is from 1:9 to 1:3.

The process of the present invention is initiated with the preparation of a solution from which the liposomes are formed. A quantity of a phosphatidyl-
5 choline, a phosphatidylglycerol and cholesterol is dissolved in an organic solvent, preferably a mixture a 1:1 (by volume) mixture of chloroform and methanol, to form a clear solution. Other solvents (and mixture thereof), such as ether, ethanol and other alcohols can be used. The preferred temperature to dissolve
10 the lipids is between room temperature and 60°C, preferably at room temperature. The solution is evaporated to form a lipid film or a lipid powder. To form a lipid film, the solvents are evaporated under nitrogen between room temperature and 60°C, preferably at room temperature. To form a lipid powder, the mixture of lipids in solution as described above is sprayed in a spray drier.
15 Preferably, the spraying takes place under nitrogen.

An aminoglycoside, for example amikacin, is dissolved in an aqueous phosphate buffer with 9% sucrose and the pH is adjusted between 6 and 8, preferably at about 7.4. The preferred buffer is 10 mM phosphate buffer. However, other aqueous buffers can be used. The aminoglycoside solution is
20 mixed with either the lipid film or the lipid powder and hydrated, preferably between 40°C and 65°C. The solution should be hydrated for at least ten minutes.

Unilamellar vesicles are formed by the application of a shearing force to the hydrated solution, e.g., by extrusion, sonication, or the use of a homogenizing apparatus such as a Gaulin homogenizer or a French press. Shearing force
25 can also be applied using injection, freezing and thawing, dialyzing away a detergent solution from lipids, or other known methods used to prepare liposomes. The size of the liposomes can be controlled using a variety of known techniques including the duration of shearing force. Preferably, the
30 modified Gaulin homogenizing apparatus described in U.S. Patent 4,753,788 is

employed to form unilamellar vesicles having diameters of less than 100 nm at a pressure of 7,000 to 13,000 psi and a temperature significantly below the transition temperature of the lipids.

The above described formulations are particularly useful for the treatment of MAC and *Pseudomonas aeruginosa* infections. The use of the above formulations indicate that MAC may be treated with significantly more amikacin delivered in a liposomal formulation than with free drug. For example it has been shown that up to 320 mg of amikacin per kilogram of mice body-weight which is more than 50% more than is tolerated with free drug. The use of the above formulations also indicates that a liposomal formulation is able to deliver significantly more amikacin to a mammal than free drug in the treatment of *P. aeruginosa*.

Amikacin delivered through a liposomal formulation is also retained in the plasma longer than free amikacin.

Since dosage regimens for aminoglycosides are well known to medical practitioners, the amount of the liposomal aminoglycoside formulation which is effective for the treatment of infections in mammals and particularly humans will be apparent to those skilled in the art.

This invention will be more fully understood by reference to the following examples, which are intended to be illustrative of the invention, but not limiting thereof. Examples 1-5 detail the formation, and both chemical and biological testing of the liposomal amikacin formulations of the invention.

Example 1

A lipid mixture of hydrogenated soy PC, cholesterol, and distearoyl phosphatidylglycerol was provided in a molar proportional of 2:1:0.1 respectively. The lipids were dissolved in chloroform and methanol (1:1 by volume). The resulting solution was stirred until the lipids dissolved and a clear solution was formed. The mixing is best carried out at room temperature. A lipid film was obtained by evaporating the organic solvents under Nitrogen at room temperature.

Amikacin free base was dissolved in 10 mM phosphate buffer with 9% sucrose and the pH adjusted to 7.4 with 6.0 M HCl. The lipid film and the amikacin solution were mixed so that the concentration of the drug was about 250 mg/ml and the concentration of the lipid was about 100 mg/ml. The resulting solution was stirred and hydrated for 10-15 minutes at 65°C. The solution was sonicated for 20 minutes using a probe sonicator (Model 500 Sonic & Material). The solution was held at 65°C for ten minutes. The solution was cooled to room temperature and centrifuged at 3600 rpm for 10 minutes. The supernatant was collected and poured through a Sephadex G-50 column to separate the liposomal formulation from the free drug. The concentration of amikacin and lipid components were determined by HPLC assay. The size was determined by optical particle sizing. The size of the liposomes varied from experiment to experiment from about 40-100 nm. For example, in one experiment a mean diameter of 62.1 nm was observed. In another, a mean diameter of 47.4 nm was observed. A preferred formulation contains a drug to lipid ratio of 1:4.

Example 2

A lipid solution was prepared in methanol and chloroform as described in Example 1. A lipid powder was obtained in a spray drier (Yamato Pulvis Basic Unit, Model GB-21). The following settings were used: 1) pump dial at 3-4.5; 2) aspirator dial at 6; 3) pressure at 1.5 -2 Kgf/cm²s; 4) inlet temperature at 50°C; and 5) outlet temperature at 40°C. The spraying took place under nitrogen for two hours. The powder was collected and combined with amikacin drug solution (as prepared in example 1). The resulting solution was mixed for 2 minutes using a high shear mixer (Virtishear) at 1000 rpm. The solution was placed in a beaker and set in a 40°C water bath and hydrated with mixing until the solution reached 40°C (about 25 minutes). The solution was then placed in a homogenizer (Gaulin 15 M) for approximately 30 passes at 10,000 psi while maintaining the inlet temperature at 40°C. The resulting solution was filtered through a 0.8 micron nylon filter. The solution was ultrafiltered to replace

unencapsulated drug with new buffer. The solution was washed with 7 to 10 volumes of buffer. The resulting product was heated to 40°C and filtered through successive 0.8, 0.45, and 0.22 micron (pore size) filters. Thus, a surprising aspect of the present invention is that the hydration of liposomes occurred significantly below the transition temperature of the formulation (about 52°C).

Example 3

The testing of liposome encapsulated amikacin for the treatment of MAC was performed using a murine model. Beige mice (C57B1/6bg^j/bg^j) were infected with MAC (101, type 1). The mice were infected by injection of 1 X 10⁷ Colony Forming Units (cfu) in the mouse tail vein (i.v.). Three experiments were performed. In the first experiment, 40, 80 and 120 milligrams of amikacin (liposomal and free) per kilogram of mouse body-weight was given i.v. daily for 5 days, beginning 7 days after infection. The animals were sacrificed 5 days after treatment was completed and the liver, lung and spleen tissue were plated. Quantitation of organisms in liver, spleen and lung tissue was performed. The cfu were determined by growth on Middlebrook 7h11 agar plates. Untreated and empty liposomes were used as controls.

The liposomes were prepared as in example 1. This experiment was performed in two identical parts. The liposomes in the first part were of an average size of 49.8 nm and the average size of the liposomes in the second part were 73.7 nm (mean diameter). The amikacin concentration was either 15.0 mg/ml or 13.21 mg/ml and the total lipid concentration was either 121 mg/ml or 55.7 mg/ml respectively (drug to lipid ratios: 0.123, 0.24). Free drug for all experiments was prepared in the same buffer that the liposomal formulations were contained in. The results are listed in Table 1. Table 1 lists the results for the spleen and liver tissue.

TABLE 1

Effects of Liposomal Amikacin on MAC Infected Mice Tissue					
Regimen	# mice	Spleen		Liver	
		cfu/g	log cfu/g	cfu/g	log cfu/g
untreated	6	1.13x10 ⁹	9.05	7.49x10 ⁸	8.87
liposomes only	3	5.30x10 ⁸	8.72	5.51x10 ⁸	8.74
amikacin 40 mg/kg	5	4.08x10 ⁸	8.61	1.41x10 ⁸	8.15
amikacin 80 mg/kg	6	4.82x10 ⁸	8.68	7.79x10 ⁷	7.89
amikacin 120 mg/kg	3	3.46x10 ⁸	8.54	5.65x10 ⁷	7.75
liposomal amikacin 40 mg/kg	6	5.43x10 ⁷	7.73	1.44x10 ⁷	7.16
liposomal amikacin 80 mg/kg	6	2.71x10 ⁷	7.43	2.11x10 ⁷	7.32
liposomal amikacin 120 mg/kg	6	4.03x10 ⁷	7.61	9.86x10 ⁶	6.99

The results for the lung study were as follows: 1) untreated regimen were 2.86 x 10⁷ cfu/gram (log = 7.46); 2) empty liposomes were 2.95 x 10⁷ cfu/gram (log = 7.47); 3) free amikacin (40 mg/kg, 80 mg/kg and 120 mg/kg) were 1.44 x 10⁶ (log = 6.16), 1.29 x 10⁶ (log = 6.11) and 9.94 x 10⁵ (log = 6.00) respectively; 4) liposomal amikacin (40 mg/kg, 80 mg/kg, and 120 mg/kg) were 2.28 x 10⁵ (log = 5.36), 3.02 x 10⁵ (log = 5.48) and 4.15 x 10⁵ (log = 5.62).

Another experiment was carried out as above (also in two parts) except that the drug therapy was started 5 days after infection. The drug was given i.v. three times/week for 21 days. The mice were sacrificed 1-2 days after the treatment stopped. During the treatment period, a group of mice received 40 mg/Kg, 80 mg/kg, and 150 mg/kg of free amikacin and another group received 40mg/kg, 80mg/kg, and 160 mg/kg of liposomal amikacin. The lung tissues were not tested. Similar controls were used as above. The liposomes were prepared as described in example 1. The average size of the liposomes was 66.5 nm or 79.6 nm (mean diameter). The concentration of the amikacin was either 15.98 mg/ml or 6.74 mg/ml. The total lipid was measure only the second part

of the experiment and it was 27.0 mg/ml (drug to lipid ratio: 1:4.167). The results are listed in Table 2.

TABLE 2

Effects of Liposomal Amikacin on MAC Infected Mice Tissue					
Regimen	# mice	Spleen cfu/g	Spleen log cfu	Liver cfu/g	Liver log cfu
untreated	6	2.23x10 ⁹	9.35	1.06x10 ⁹	9.03
liposomes only	5	2.27x10 ⁹	9.36	2.08x10 ⁹	9.32
amikacin 40 mg/kg	6	5.76x10 ⁸	8.76	1.09x10 ⁸	8.04
amikacin 80 mg/kg	6	2.02x10 ⁸	8.31	3.67x10 ⁷	7.56
amikacin 120 mg/kg	2	1.48x10 ⁸	8.17	3.15x10 ⁷	7.50
liposomal amikacin 40 mg/kg	6	7.98x10 ⁷	7.90	7.24x10 ⁶	6.86
liposomal amikacin 80 mg/kg	4	2.57x10 ⁷	7.41	2.21x10 ⁶	6.34
liposomal amikacin 160 mg/kg	6	1.73x10 ⁷	7.24	1.04x10 ⁶	6.02

A third experiment was carried out as in the second experiment except that the mice treated with the liposomal amikacin received 120 mg/kg, 240 mg/kg and 320 mg/kg doses and the mice treated with the free amikacin received only 120 mg/kg doses. The liposomes were prepared as in example 2 where the average size of the liposomes were 81.8 nm (median diameter). The amikacin concentration was 32.02 mg/ml and the total lipid concentration was 91.5 mg/ml (drug to lipid ratio: 1:2.941). The results are listed in Table 3.

TABLE 3

Effects of Liposomal Amikacin on MAC Infected Mice Tissue					
Regimen	# mice	Spleen		Liver	
		cfu/g	log cfu/g	cfu/g	log cfu/g
untreated	6	3.70x10 ⁹	9.57	1.65x10 ⁹	9.22
liposomes only	6	4.05x10 ⁸	8.61	1.20x10 ⁸	8.08
amikacin 120 mg/kg	6	2.13x10 ⁸	8.33	6.10x10 ⁷	7.78
liposomal amikacin 120 mg/kg	6	8.8x10 ⁶	6.94	3.67x10 ⁵	5.56
liposomal amikacin 240 mg/kg	6	5.52x10 ⁶	6.74	4.83x10 ⁵	5.68
liposomal amikacin 320 mg/kg	6	<3.38x10 ⁶	6.53	<2.00x10 ⁵	5.30

The results of all the experiments establish that significantly higher doses of amikacin can be delivered without an increase in toxicity and with superior efficacy. It is known in the art that 150 mg/kg of amikacin will kill many of the mice injected. Thus, the delivery of 320 mg/kg of amikacin is a significant increase in the amount of drug delivered without lethal toxicity.

Example 4

The efficacy and toxicity of liposomal amikacin in *Pseudomonas* infected mice was tested. CF-1 mice were used (females, 6-8 weeks old). The mice were obtained from Jackson Labs. A clinical isolate of *Pseudomonas aeruginosa* was obtained on a Mueller Hinton/MacConkey blood agar plate. The organisms were transferred to a Mueller Hinton plate and grown for 24 hours. Colonies were transferred to saline and grown for 48 hours. The samples were frozen in saline containing 15% fetal calf serum at 1 x 10⁸ organisms/ml (MacFarland Standard). The colonies were stored at -70°C. The colonies were thawed and grown in a Mueller Hinton plate for 24 hours. The bacteria were adjusted to 8 x 10⁶ organisms/ml in saline containing talc (62.5 mg/ml) and 1 ml was injected intraperitoneal. A culture of inoculum on MH agar for 24 hours determined that approximately 7 x 10⁶ cfu were delivered per mouse. Liposomal amikacin and empty liposomes were prepared as set out in example 2. The liposomes of the

amikacin formulation had an average size (median diameter) of 62.4 nanometers. The lipid concentration was 95.31 mg/ml and the amikacin concentration was 23.54 mg/ml (.25 drug/lipid ratio). The pH of the solution containing the formulation was 7.31. The empty liposomes had an average size of 66.8
 5 nanometers with 92.05 mg/ml of lipid. The concentration of the free amikacin solution was 23.54 mg/ml. The pH of the solution containing the liposomes was 7.33.

Mice were treated with 40 mg/kg (drug per body-weight), 80 mg/kg, 120 mg/kg and 240 mg/kg of liposomal amikacin. Mice were also treated with 40
 10 mg/kg, 80 mg/kg, and 120 mg/kg of free amikacin. The drugs were administered intravenously in the caudal vein. Two doses were given; one at four hours after infection and one at twenty-four hours after infection. The results are listed in Table 4. The results establish that it is possible to deliver up to 240 mg/kg of amikacin using a liposome formulation with all the subjects surviving.

15

TABLE 4

20

25

Effects of Liposomal Amikacin on Pseudomonas Infection in Mice		Mortality #died/#tested hours post infection				
Infection	Treatment	24	30	48	72	%
no	none	0/4	0/4	0/4		0
yes	none	5/6		6/6		100
yes	40 (free amikacin)	1/4	1/4	1/4	1/4	25
yes	80 (free amikacin)	0/6	0/6	0/6	0/6	0
yes	120 (free amikacin)	0/6	0/6	0/6	0/6	0
yes	40 (Lipo. amikacin)	1/4	3/4	4/4	4/4	100
yes	80 (lipo. amikacin)	0/6	0/6	1/6	2/6	33
yes	120 (lipo. amikacin)	0/6	0/6	0/6	0/6	0
yes	240 (lipo. amikacin)	0/6	0/6	0/6	0/6	0

Example 5

Amikacin content was measured in mouse (C57Bl/6, females 2-3 months old) plasma after injection (i.v.) of free amikacin or liposomal amikacin (100 mg/kg). Liposomal amikacin was prepared as described in example 2. Blood samples were obtained at various time intervals. The samples were collected by removing 100 microliters of blood, retroorbitally, from nonanesthetized mice. The samples were collected in heparinized capillary pipets which were plugged with modeling clay. The samples were centrifuged at 3250 rpm (20 cm rotor) for 10 minutes. The samples were analyzed using a radioimmunoassay (Diagnostic Products). The liposomal amikacin formulation contained liposomes with an average size of 43.6 nm (median diameter). The amikacin concentration was 10.09 mg/ml and the total lipid concentration was 56.8 mg/ml (drug to lipid ratio 1 : 5.556). The results are listed in Table 5.

TABLE 5

15

Amikacin Content in Mouse Plasma				
Treatment	Interval (hours)	# of mice	Average Concentration ($\mu\text{g/ml}$)	Standard deviation ($\pm\mu\text{g/ml}$)
Amikacin	0.0833	4	406	155
	0.25	4	106	12
	2	4	65	12
	6	4	none detected	
	14		not tested	
	24		not tested	
	Liposomal Amikacin	0.0833	4	530
0.25		4	344	321
2		3	346	199
6		4	278	137
14		4	272	78
24		6	108	14

20

The results establish that liposomal amikacin is retained in the plasma for a significantly longer period than free amikacin.

Although this specification has been disclosed and illustrated with reference to particular applications, the principles involved are susceptible to numerous other applications which will be apparent to those skilled in the art. The invention is, therefore, to be limited only as indicated by the scope of the appended claims.

What is claimed is:

1. Liposomes comprising an encapsulated aminoglycoside wherein the liposomes are unilamellar vesicles having an average size of less than 100 nm which are stable in liquid form for at least 2 weeks at 22°C without change in size or loss of more than 10% of the encapsulated aminoglycoside.
2. The formulation of claim 1 wherein the liposomes are comprised of a neutral lipid and a sterol.
3. The formulation of claim 2 wherein the liposomes are further comprised of a negatively charged lipid.
4. The formulation of claims 1, 2, or 3 wherein the molar amount of negatively charged is less than 20% total lipid.
5. The formulation of claim 3 in which the liposomes are comprised of phosphatidylcholine, cholesterol and phosphatidylglycerol, wherein the molar amount of phosphatidylglycerol is less than 20% of total molar lipid.
6. The formulation of claim 3 in which the liposomes are comprised of hydrogenated soy phosphatidylcholine, cholesterol and distearoyl phosphatidylglycerol in a molar ratio of about 2:1:0.1, wherein the aminoglycoside to total lipid ratio is between 1:9 to 1:3.
7. A process for making aminoglycoside liposomes comprising:
 - (a) forming a lipid powder comprised of a neutral phospholipid and a sterol;
 - (b) mixing the powder with an aminoglycoside in a buffer;
 - (c) hydrating the mixture at a temperature significantly below the transition temperature of the lipid mixture;
 - (d) reducing the size of the liposomes to an average size of less than 100 nm by the application of a shearing force.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 92/10591

A. CLASSIFICATION OF SUBJECT MATTER

IPC5: A61K 9/127, A61K 31/71
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC5: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EMBASE, CA, WPI, WPIL, CLAIMS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4753788 (R C GAMBLE), 28 June 1988 (28.06.88), column 2, line 21 - line 59, claims --	1-7
Y	WO, A1, 8804573 (THE LIPSOME COMPANY, INC.), 30 June 1988 (30.06.88), claims 1-61 --	1-7

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "B" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art


"&" document member of the same patent family

Date of the actual completion of the international search

6 August 1993

Date of mailing of the international search report

09.09.93

Name and mailing address of the ISA/
 European Patent Office, P.B. 5818 Patendaan 2
 NL-2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo rd,
 Fax: (+31-70) 340-3016

Authorized officer

ANNELI JÖNSSON

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 92/10591

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Dialog Information Service, EMBASE, Dialog accession no. 8333576/5, EMBASE no. 92010722, Wichert B.V. et al: "Amikacin liposomes: Characterization, aerosolization, and in vitro activity against Mycobacterium avium-intracellulare in alveolar macrophages", INT. J. PHARM. (Netherlands), 1992, 78/2-3 (227-235) ----- -----	1-7

SA 3012

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/07/93

International application No.
PCT/US 92/10591

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4753788	28/06/88	CA-A- 1282699 EP-A,B- 0190050 JP-A- 61218515	09/04/91 06/08/86 29/09/86
WO-A1- 8804573	30/06/88	AU-B- 609711 AU-A- 6934587 CA-A- 1314481 DE-A- 3785198 EP-A,B- 0295248 EP-A- 0498471 EP-A- 0500143 US-A- 4981692	09/05/91 15/07/88 16/03/93 06/05/93 21/12/88 12/08/92 26/08/92 01/01/91